



**CALIFORNIA SCIENCE & ENGINEERING FAIR
2018 PROJECT SUMMARY**

Name(s) <p align="center">Danya Balagopal</p>	Project Number <p align="right">38338</p>
Project Title <p align="center">Constructing a Sustainable, Low-Cost Herbal Biosorbents Filter to Remove Heavy Metals from Contaminated Groundwater</p>	
<p align="center">Abstract</p> <p>Objectives/Goals 22% of California's community water systems rely on contaminated groundwater. Activated carbon filters are expensive and nonrenewable. The goal of my project is to construct a sustainable, low-cost filter by investigating the biosorptive effect of torrefied <i>Oryza sativa</i> hull pellets, <i>Moringa oleifera</i> seed kernels, <i>Vetiveria zizanioides</i> roots, <i>Azadirachta indica</i> leaves, and the adsorptive effect of Kaolinite, and Kaolinite-<i>Carica papaya</i> on the removal of iron and copper by 75% and reduce turbidity by 50%.</p> <p>Methods/Materials Torrefied <i>O. sativa</i> hulls, <i>M. oleifera</i> seed, <i>V. zizanioides</i> roots, <i>A. indica</i> leaves, kaolinite and kaolinite-<i>C. papaya</i> clay bowls were tested by soaking each of them in groundwater and varying: Temperature (100,110,120C) Adsorbent dose (5g,10g,15g) Contact times (60,120,180 minutes) with four trials each against the control of untreated groundwater. A Sper Scientific Turbidity meter was used to test turbidity. The removal efficiency was calculated and analyzed through ANOVA and supported by literature.</p> <p>Results My experiments supported the hypotheses proving that the biosorbents could remove iron and copper from groundwater by more than 75%. All tested biosorbents removed iron by 100%. <i>V. zizanioides</i> and the Kaolinite hybrid clay removed copper by 100%, while the others were at 80%. <i>M. oleifera</i>, <i>V. zizanioides</i>, and Kaolinite reduced turbidity by 50% and Kaolinite hybrid clays by 70%. However, <i>O. sativa</i> and <i>A. indica</i> did not support the hypothesis as they increased turbidity by 105% and 80% respectively.</p> <p>Conclusions/Discussion All the biosorbents adsorbed metals for different reasons. Torrefying <i>O. sativa</i> created a porous cell structure that increased the surface sites available for metal ion adsorption. The herbs adsorbed through complexation although they contain different compounds: both <i>M. oleifera</i>, <i>V. zizanioides</i> contain saponin, while <i>A. indica</i> contain salannin and azidirectin. Kaolinite clay possesses a high ion exchange capacity, while kaolinite-<i>C. papaya</i> has a high cation exchange capacity. <i>M. oleifera</i>, kaolinite and kaolinite-<i>C. papaya</i> possess strong flocculation and/or coagulative properties which reduced turbidity. My product which consists of a kaolinite- <i>C. papaya</i> hybrid bowl with <i>V. zizanioides</i> roots and a water-soluble capsule containing <i>M. oleifera</i> only costs \$1.35 to produce. This provides a low-cost and sustainable alternative to carbon filters for rural communities dependent on groundwater.</p>	
Summary Statement My project investigated the effect of six biosorbents on the removal of iron and copper and turbidity reduction in groundwater, to construct a sustainable, low-cost herbal filter as an alternative to activated carbon filters.	
Help Received I designed, experimented, and analyzed my results on my own. I thank my teacher, Dr. Wilmot, for his support; Mr. Dan Coltrin (Forensic Laboratories) for answering my questions; Clay Planet for firing my kaolinite bowls.	